

GenCore version 5.1.4\_p5\_4578  
Copyright (c) 1993 - 2003 CompuGen Ltd.

OM protein - protein search, using sw model

Run on: May 19, 2003, 16:35:42 ; Search time 53.4605 Seconds  
(without alignments)  
897.302 Million cell updates/sec

Title: US-09-625-573-4  
Perfect score: 1900  
Sequence: 1-MLSTSRSEIRNFNSGEV.....DGVTSNTPSTGCEQVSAVL 360

Scoring table: BLOSUM62  
Gapop 10.0 , Gapext 0.5  
Searched: 908470 seqs, 133250620 residues  
Total number of hits satisfying chosen parameters: 908470

Minimum DB seq length: 0  
Maximum DB seq length: 2000000000

Post-processing: Minimum Match 0%  
Maximum Match 100%  
Listing first 45 summaries

Database :			
A.Geneseq.101002:*			
1:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1980.DAT:*		
2:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1981.DAT:*		
3:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1982.DAT:*		
4:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1983.DAT:*		
5:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1984.DAT:*		
6:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1985.DAT:*		
7:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1986.DAT:*		
8:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1987.DAT:*		
9:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1988.DAT:*		
10:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1989.DAT:*		
11:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1990.DAT:*		
12:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1991.DAT:*		
13:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1992.DAT:*		
14:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1993.DAT:*		
15:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1994.DAT:*		
16:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1995.DAT:*		
17:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1996.DAT:*		
18:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1997.DAT:*		
19:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1998.DAT:*		
20:	/SID22/gcgdata/geneseq/geneseq-emb1/AA1999.DAT:*		
21:	/SID22/gcgdata/geneseq/geneseq-emb1/AA2000.DAT:*		
22:	/SID22/gcgdata/geneseq/geneseq-emb1/AA2001.DAT:*		
23:	/SID22/gcgdata/geneseq/geneseq-emb1/AA2002.DAT:*		

Pred. No. is the number of results predicted by chance to have a score greater than or equal to the score of the result being printed, and is derived by analysis of the total score distribution.

SUMMARIES

Result No.	Query Score	Match Length	ID	Description
1	1900	100.0	16	AA79166 Human monocyte che
2	1900	100.0	18	AAW35833 Human monocyte che
3	1900	100.0	22	AAW80108 Human CCR2b protei
4	1900	100.0	360	AAW07614 Human wild-type CC
5	1899	99.9	360	AAU07613 Human CCR2-641 pol
6	1894	99.7	360	AAU07613 Human CCR2-641 pol
7	1651	86.9	374	16 AAB56340 Non-endogenous hum
8	1651.5	86.9	374	22 AAG80107 Human monocyte-che
9	1473	77.5	329	22 AAB46859 Human MCP-1 recept
10	1386	72.9	19	AAW54037 Mouse CC-CR5 prot

11	1371	72.2	352	22	AAG79089	Amino acid sequenc
12	1364	71.8	352	18	AAW27407	Human CCR5. Homo
13	1364	71.8	352	18	AAW27123	Human chemokine re
14	1364	71.8	352	19	AAW27125	Macaque chemokine
15	1364	71.8	352	19	AAW23835	Human CC chemokine
16	1364	71.8	352	20	AAW88232	HIV-1 co-receptor
17	1364	71.8	352	22	AAW80111	Human CCR5 protein
18	1364	71.8	352	22	AAW82948	Human HIV-1 co-rec
19	1364	71.8	352	22	AAW83354	Human CCR5 protein
20	1364	71.8	352	22	AAE04321	Human chemokine re
21	1364	71.8	352	23	ABB08343	Human chemokine (C
22	1364	71.8	352	23	AAW52828	Human CC chemokine
23	1364	71.8	439	20	AAW41280	Fusion protein con
24	1359.5	71.6	371	19	AAW23834	Human CC chemokine
25	1358	71.5	352	22	ABB56342	Non-endogenous hum
26	1358	71.5	352	23	AAW52829	Human CCR5 Gln 55
27	1356	71.4	352	22	AAE07039	Human G-protein ch
28	1356	71.4	352	22	AAE07048	Human G-protein ch
29	1356	71.4	352	22	AAW64858	Human HGRK10 prot
30	1356	71.4	352	23	AAW097152	Human G-protein ch
31	1355	71.3	352	18	AAW07602	Human G-protein ch
32	1355	71.3	352	21	AAW80128	Human G-protein ch
33	1355	71.3	352	22	AAE07037	Human G-protein ch
34	1355	71.3	352	22	AAE07046	Human G-protein ch
35	1355	71.3	352	23	AAW097150	Human chemokine re
36	1335.5	70.3	332	18	AAW26766	C-C chemokine rece
37	1036	54.5	355	15	AAW52749	Human MIP-1 alpha/
38	1036	54.5	355	18	AAW26588	Human MIP-1alpha/R
39	1036	54.5	355	21	AAW25751	Human CC-chemokine
40	1036	54.5	355	18	AAW20571	Rat CC chemokine r
41	995.5	52.4	355	18	AAW29179	Human C-C chemokin
42	958	50.4	355	19	AAW51744	CC-chemokine recep
43	951	50.1	355	17	AAW03376	Human C-C chemokin
44	951	50.1	355	18	AAW10300	Human C-C chemokin
45	951	50.1	355	23	ABB07733	Human C-C chemokin

ALIGNMENTS

RESULT 1  
AA79166  
ID AA79166 standard; Protein; 360 AA.

XX AA79166;

XX 29-DEC-1995 (first entry)

XX Human monocyte chemoattractant protein-1 receptor MCP-1RB.

XX Monocyte chemoattractant protein-1 receptor; MCP-1R; chemokine.

XX Homo sapiens.

XX	Key	Location/Qualifiers
FT	Domain	49..70
FT	Domain	/label= transmembrane
FT	Domain	80..700
FT	Domain	/label= transmembrane
FT	Domain	115..136
FT	Domain	/label= transmembrane
FT	Domain	154..178
FT	Domain	/label= transmembrane
FT	Domain	204..231
FT	Domain	/label= transmembrane
FT	Domain	244..268
FT	Domain	/label= transmembrane
FT	Domain	295..313
FT	Domain	/label= transmembrane
FT	Region	314..360
FT	Domain	/label= carboxyl tail
FT	Domain	1..48
FT	Domain	/label= extracellular

XX WO9519436-A.  
 XX 20-JUL-1995.  
 XX 11-JAN-1995; 95WO-US00476.  
 XX 13-JAN-1994; 94US-0182962.  
 XX (REGC ) UNIV CALIFORNIA.  
 XX Charo I, Coughlin S;  
 XX WPI; 1995-263866/34.  
 XX N-PSDB; AAQ96298.  
 XX  
 XX DNA encoding monocyte chemo-attractant protein-1 receptor - used partic.  
 XX for identifying antagonists and for treating diseases characterised by  
 XX monocyte infiltrates

Claim 2; Fig 2; 84pp; English.

To identify and clone new members of the chemokine receptor gene family, degenerate oligo primers were designed corresp. to the conserved sequences R79167 in the second and R79168 in the third transmembrane domains of the MIP-1alpha/RANTES receptor, the IL-8 receptors and the HUMSRS orphan receptor (GenBank Accession #M99293). The degenerate oligo incorporating EcoRI and XhoI sites at their 5' ends are Q96299 and Q96300. Amplification of cDNA derived from MM6 cells with the primers yielded a number of PCR products. One cDNA appeared to encode a novel protein. To obtain a full-length version of this clone, a MM6 cDNA library was constructed in pFRG and probed with the PCR product. A 2.1 kb cDNA clone was obtained. Analysis of additional clones in the MM6 cDNA library revealed a second sequence that was identical to the 2.1 kb cDNA sequence first obtained from the 5' UTR through the putative seventh transmembrane domain but contained a different cytoplasmic tail. The second sequence appears to represent alternative splicing of the carboxyl-terminal tail of the MCP-1R protein. The two sequences are denoted MCP-1RA and MCP-1RB (see Q96297/R79165 & Q96298/R79166). Active mature MCP-1RA has a predicted mol. wt. of about 42,000 daltons. MCP-1RB has a mol. wt. of about 41,000 daltons.

XX Sequence 360 AA;

Query Match 100.0%; Score 1900; DB 16; Length 360;  
 Best Local Similarity 100.0%; Pred. No. 3.7e-211;  
 Matches 360; Conservative 0; Mismatches 0; Indels 0; Gaps 0;  
 1 MLSTSRSFIRNTNSESGETTFFDYDYGAPCHKFDVKQIGAOQLPPLYSLVFIQFVGN 60  
 1 MLSTSRSFIRNTNSESGETTFFDYDYGAPCHKFDVKQIGAOQLPPLYSLVFIQFVGN 60  
 61 MLVVLILINCKKLCITDIYLLNLAISDILLPLPLWAHSAANWVFGNCKLFTGLY 120  
 61 MLVVLILINCKKLCITDIYLLNLAISDILLPLPLWAHSAANWVFGNCKLFTGLY 120  
 121 HIGYFGGIFFIILLTDRLAIVHAFKARTVTVGVVTSVITLWVAFASVPGIIFTK 180  
 121 HIGYFGGIFFIILLTDRLAIVHAFKARTVTVGVVTSVITLWVAFASVPGIIFTK 180  
 181 COKEDSVVCGPYFPRGWNNEFTIMRNILGLVPLLMVICYSGILKTLRCRNEKKRHR 240  
 181 COKEDSVVCGPYFPRGWNNEFTIMRNILGLVPLLMVICYSGILKTLRCRNEKKRHR 240  
 241 AVRVIITMIVYFLFWTPYNIIVLLNTQEFGLSNCESTSLDQATQVETILGMTHCCI 300  
 241 AVRVIITMIVYFLFWTPYNIIVLLNTQEFGLSNCESTSLDQATQVETILGMTHCCI 300  
 301 NPILYAFVGEKFRRLYSVFFRKHITKRCCKQCPVYRETVDGVTSTNTPTSTGEQVYSAGL 360  
 301 NPILYAFVGEKFRRLYSVFFRKHITKRCCKQCPVYRETVDGVTSTNTPTSTGEQVYSAGL 360

XX RESULT 3

RESULT 2

AAW35833  
 ID AAW35833 standard; Protein; 360 AA.

XX AAW35833;

XX 27-FEB-1998 (first entry)

XX Human monocyte chemoattractant protein 1 receptor.

XX Human; MCP-1; monocyte chemoattractant protein; receptor; tumour;  
 XX inflammatory disease; viral; allergy; diabetes.

XX Homo sapiens.

XX JP09238688-A.

XX 16-SEP-1997.

XX 11-MAR-1996; 96JP-0053574.

XX 11-MAR-1996; 96JP-0053574.

XX (TAKE ) TAKEDA CHEM IND LTD.

XX WPI; 1997-506557/47.

XX N-PSDB; AAT96976.

XX DNA encoding human monocyte chemoattractant protein 1 receptor -  
 XX used to treat tumours and inflammatory, viral, infectious, allergic,  
 XX diabetic and central nervous system diseases

XX Disclosure; Page 12-14; 15pp; Japanese.

XX The present sequence represents human monocyte chemoattractant protein 1  
 XX (MCP-1) receptor protein. The MCP-1 receptor protein and encoding DNA  
 XX are used for the prevention and treatment of tumours and inflammatory,  
 XX viral, infectious, allergic, diabetic and central nervous system  
 XX diseases.

XX Sequence 360 AA;

Query Match 100.0%; Score 1900; DB 18; Length 360;  
 Best Local Similarity 100.0%; Pred. No. 3.7e-211;  
 Matches 360; Conservative 0; Mismatches 0; Indels 0; Gaps 0;  
 1 MLSTSRSFIRNTNSESGETTFFDYDYGAPCHKFDVKQIGAOQLPPLYSLVFIQFVGN 60  
 1 MLSTSRSFIRNTNSESGETTFFDYDYGAPCHKFDVKQIGAOQLPPLYSLVFIQFVGN 60  
 61 MLVVLILINCKKLCITDIYLLNLAISDILLPLPLWAHSAANWVFGNCKLFTGLY 120  
 61 MLVVLILINCKKLCITDIYLLNLAISDILLPLPLWAHSAANWVFGNCKLFTGLY 120  
 121 HIGYFGGIFFIILLTDRLAIVHAFKARTVTVGVVTSVITLWVAFASVPGIIFTK 180  
 121 HIGYFGGIFFIILLTDRLAIVHAFKARTVTVGVVTSVITLWVAFASVPGIIFTK 180  
 181 COKEDSVVCGPYFPRGWNNEFTIMRNILGLVPLLMVICYSGILKTLRCRNEKKRHR 240  
 181 COKEDSVVCGPYFPRGWNNEFTIMRNILGLVPLLMVICYSGILKTLRCRNEKKRHR 240  
 241 AVRVIITMIVYFLFWTPYNIIVLLNTQEFGLSNCESTSLDQATQVETILGMTHCCI 300  
 241 AVRVIITMIVYFLFWTPYNIIVLLNTQEFGLSNCESTSLDQATQVETILGMTHCCI 300  
 301 NPILYAFVGEKFRRLYSVFFRKHITKRCCKQCPVYRETVDGVTSTNTPTSTGEQVYSAGL 360  
 301 NPILYAFVGEKFRRLYSVFFRKHITKRCCKQCPVYRETVDGVTSTNTPTSTGEQVYSAGL 360

XX RESULT 3

AA80108  
ID AAG80108 standard; Protein: 360 AA.  
XX AAG80108;  
XX 17-JAN-2002 (first entry)  
XX Human CCR2b protein.  
XX Chemokine; tumour diagnosis; colorectal; prostatic; organ rejection;  
KW inflammation; autoimmune disease; metastasis; bronchial asthma; lupus;  
KW chronic bowel inflammation; rheumatoid arthritis; cytostatic;  
KW antiinflammatory; antiasthmatic; immunosuppressive; dermatological;  
KW antirheumatic; antiarthritic.  
XX Homo sapiens.  
OS  
XX WO200172830-A2.  
XX 04-OCT-2001.  
XX 02-APR-2001; 2001WO-EP03708.  
XX 31-MAR-2000; 2000DE-1016013.  
XX (IPFP-) IPF PHARM GMBH.  
XX (FORS/) FORSMANN U.  
XX Forssmann W, Adermann K, Heitland A, Spodsborg N;  
XX WPI; 2001-626256/72.  
XX Diagnostic agent containing two or more receptor-specific ligands,  
PT useful for detecting tumors, inflammation etc., also therapeutic use of  
PT ligand inhibitors  
XX  
XX Disclosure; Page 9; 26pp; German.  
XX  
XX This invention describes a novel diagnostic agent (A) comprising at least  
CC two different ligands (I) for receptors (II) that are implicated in  
CC disease. (A) are used for the diagnosis of tumors (especially colorectal  
CC or prostatic), organ rejection, inflammation and autoimmune diseases.  
CC Also inhibitors of (I) are used therapeutically against tumors (and their  
CC metastases), inflammation (particularly bronchial asthma or chronic bowel  
CC inflammation), or autoimmune diseases (rheumatoid arthritis or lupus).  
CC where the (cardio)vascular, lymphatic, respiratory, nervous, digestive,  
CC endocrine, motor or urogenital systems or skin are affected, and bone  
CC marrow diseases. The products of the invention are chemokine derivatives  
CC which have cytostatic, antiinflammatory, antiasthmatic,  
CC immunosuppressive, dermatological, antirheumatic, antiarthritic.  
CC Chemokines act on specific tumor and inflammatory cells through a  
CC constellation of chemokine receptors (CR), which control migration and  
CC proliferation of these cells. AAG8045-AAG80128 represent human chemokine  
CC fragments used to illustrate the method of the invention.  
XX  
XX Sequence 360 AA;  
Query Match 100.0%; Score 1900; DB 22; Length 360;  
Best Local Similarity 100.0%; Pred. No. 3.7e-211;  
Matches 360; Conservative 0; Mismatches 0; Indels 0; Gaps 0;  
QY 1 MLSTSRSRFRINTNESGEVTTFFDYDGAPCHKFDVKQIGQALLPPLYSLVFIFGVGN 60  
DB 1 MLSTSRSRFRINTNESGEVTTFFDYDGAPCHKFDVKQIGQALLPPLYSLVFIFGVGN 60  
QY 61 MLVLLINLCKLCLTDIYLLNLALISDLLFLITLPLWAHSAANWVFCNAMCKLFTGLY 120  
DB 61 MLVLLINLCKLCLTDIYLLNLALISDLLFLITLPLWAHSAANWVFCNAMCKLFTGLY 120  
QY 121 HIGVFGGIFILLIDIRYLAIVHAFKARTVTGCVVTSVITWLVAVFASVPGIIFTK 180  
DB 121 HIGVFGGIFILLIDIRYLAIVHAFKARTVTGCVVTSVITWLVAVFASVPGIIFTK 180

QY 181 CQKEDSVYVCGPYFPRGNNFHTIMRNLTGLVLPLLMVICYSGILKTLKCRNEKKRHR 240  
DB 181 CQKEDSVYVCGPYFPRGNNFHTIMRNLTGLVLPLLMVICYSGILKTLKCRNEKKRHR 240  
QY 241 AVRVIETIMIVYELFWTPYINIVILLNTFQEFFGLSNCESTSQDQATQVETLGMTHCCI 300  
DB 241 AVRVIETIMIVYELFWTPYINIVILLNTFQEFFGLSNCESTSQDQATQVETLGMTHCCI 300  
QY 301 NPIIYAFVGGKFRYLSVFFRKHITKRFCKQCPVYRETVDGVTSTNTPTSGEQEVSAGL 360  
DB 301 NPIIYAFVGGKFRYLSVFFRKHITKRFCKQCPVYRETVDGVTSTNTPTSGEQEVSAGL 360  
RESULT 4  
AAU07614  
ID AAU07614 standard; Protein: 360 AA.  
XX AAU07614;  
XX 04-DEC-2001 (first entry)  
XX Human wild-type CCR2-64V polypeptide.  
XX Human; CCR2 receptor; CCR2-64I; CCR2-64V; gene therapy; atherosclerosis;  
KW single nucleotide polymorphism; hypercholesterolaemia.  
XX Homo sapiens.  
XX WO200162796-A1.  
XX 30-AUG-2001.  
XX 22-FEB-2001; 2001WO-GB00755.  
XX 22-FEB-2000; 2000GB-0004183.  
XX (SMIK ) SMITHKLINE BEECHAM PLC.  
XX Valdes AM, Groot PHE, Spurr NK;  
XX WPI; 2001-550086/61.  
XX N-PSDB; AAS12140.  
XX Diagnosing atherosclerosis or susceptibility to atherosclerosis in a  
PT subject, by determining a single nucleotide polymorphism in specific  
PT codon of a polynucleotide encoding human CCR2 receptor in genome of the  
PT subject  
XX Claim 1; Page 21; 28pp; English.  
XX The invention relates to diagnosing atherosclerosis (or susceptibility  
CC to) in a subject by determining expression or activity of the human  
CC CCR2-64I polypeptide (a polymorphic variant form of the human CCR2  
CC receptor) or the CCR2-64V polypeptide (human CCR2 receptor), by screening  
CC for a single nucleotide polymorphism in codon 64 of the polynucleotide  
CC encoding the CCR2 receptor. This results in production of CCR2-64I,  
CC whereby polymorphic variants are associated with a lower incidence of  
CC atherosclerosis. The presence or amount of CCR2-64I/V in a sample can  
CC also be analysed. The sequences of the invention can be used for  
CC predicting the response of a patient to drug treatment, for predicting  
CC the disease outcome in a patient and also for the production of a  
CC treatment for hypercholesterolaemia. The sequence represents the  
CC wild-type receptor polypeptide CCR2-64V.  
XX  
XX Sequence 360 AA;  
Query Match 100.0%; Score 1900; DB 22; Length 360;  
Best Local Similarity 100.0%; Pred. No. 3.7e-211;  
Matches 360; Conservative 0; Mismatches 0; Indels 0; Gaps 0;  
QY 1 MLSTSRSRFRINTNESGEVTTFFDYDGAPCHKFDVKQIGQALLPPLYSLVFIFGVGN 60  
DB 1 MLSTSRSRFRINTNESGEVTTFFDYDGAPCHKFDVKQIGQALLPPLYSLVFIFGVGN 60



QY 61 MLVVLILNCKKLCITDIYLLNLAISDLLFLITPLWAHSAANEVFGNAMCKLFTGLY 120  
 DB 61 MLVVLILNCKKLCITDIYLLNLAISDLLFLITPLWAHSAANEVFGNAMCKLFTGLY 120  
 QY 121 HIGYFGGIFILLITDRYLAIHVAHFALKARTVTFGVVTSVITLWVAFASVPGIIFTK 180  
 DB 121 HIGYFGGIFILLITDRYLAIHVAHFALKARTVTFGVVTSVITLWVAFASVPGIIFTK 180  
 QY 181 COKEDSVYVCGPYPRGWNFFHIMRNILGLVLPPLIMVICYSGILKTLRCRNEKKRHR 240  
 DB 181 COKEDSVYVCGPYPRGWNFFHIMRNILGLVLPPLIMVICYSGILKTLRCRNEKKRHR 240  
 QY 241 AVRVIPTIMIVFLFPTPNVIVLLNTFQEFFGLSNCESTSQLDQAVQVTTETLGMTHCCI 300  
 DB 241 AVRVIPTIMIVFLFPTPNVIVLLNTFQEFFGLSNCESTSQLDQAVQVTTETLGMTHCCI 300  
 QY 301 NPIIYAFVGEKFRRLSVFRRKHITRFRCKQCPVFYRETVDGVTSTNTPSTGEOEVSAGL 360  
 DB 301 NPIIYAFVGEKFRRLSVFRRKHITRFRCKQCPVFYRETVDGVTSTNTPSTGEOEVSAGL 360

## RESULT 5

AAU07613  
 ID AAU07613 standard; Protein; 360 AA.

AC AAU07613;

DT 04-DEC-2001 (first entry)

DE Human CCR2-64I polymorphic variant polypeptide.

KW Human; CCR2 receptor; CCR2-64I; CCR2-64V; gene therapy; atherosclerosis;  
 KW single nucleotide polymorphism; hypercholesterolaemia.

OS Homo sapiens.

FH Key Location/Qualifiers

FT Misc-difference 64 /note= "wild-type Val is replaced by Ile"

PN WO200162796-A1.

PD 30-AUG-2001.

PF 22-FEB-2001; 2001WO-GB00755.

PR 22-FEB-2000; 2000GB-0004183.

PS (SMK ) SMITHKLINE BEECHAM PLC.

PI Valdes AM, Groot PHE, Spurr NK;

DR WPI; 2001-550086/61.

DR N-PSDB; AAS12139.

PT Diagnosing atherosclerosis or susceptibility to atherosclerosis in a  
 PT subject, by determining a single nucleotide polymorphism in specific  
 PT codon of a polynucleotide encoding human CCR2 receptor in genome of the  
 PT subject.

PS Claim 1; Page 20; 28pp; English.

XX The invention relates to diagnosing atherosclerosis (or susceptibility  
 CC to) in a subject by determining expression or activity of the human  
 CC CCR2-64I polypeptide (a polymorphic variant form of the human CCR2  
 CC receptor) or the CCR2-64V polypeptide (human CCR2 receptor), by screening  
 CC for a single nucleotide polymorphism in codon 64 of the polynucleotide  
 CC encoding the CCR2 receptor. This results in production of CCR2-64I,  
 CC whereby polymorphic variants are associated with a lower incidence of  
 CC atherosclerosis. The presence or amount of CCR2-64I/V in a sample can  
 CC also be analysed. The sequences of the invention can be used for  
 CC predicting the response of a patient to drug treatment, for predicting

CC the disease outcome in a patient and also for the production of a  
 CC treatment for hypercholesterolaemia. The sequence represents the  
 CC polymorphic variant polypeptide CCR2-64I.

SQ Sequence 360 AA;

Query Match 99.9%; Score 1899; DB 22; Length 360;

Best Local Similarity 99.7%; Pred. No. 4.8e-211;

Matches 359; Conservative 1; Mismatches 0; Indels 0; Gaps 0;

QY 1 MLSTSRSRFIRNTNSESGETVTFDDYDYGAPCHKFDVKQIGAQLLPPLYSLVIFGFGVGN 60

DB 1 MLSTSRSRFIRNTNSESGETVTFDDYDYGAPCHKFDVKQIGAQLLPPLYSLVIFGFGVGN 60

QY 61 MLVVLILNCKKLCITDIYLLNLAISDLLFLITPLWAHSAANEVFGNAMCKLFTGLY 120

DB 61 MLVVLILNCKKLCITDIYLLNLAISDLLFLITPLWAHSAANEVFGNAMCKLFTGLY 120

QY 121 HIGYFGGIFILLITDRYLAIHVAHFALKARTVTFGVVTSVITLWVAFASVPGIIFTK 180

DB 121 HIGYFGGIFILLITDRYLAIHVAHFALKARTVTFGVVTSVITLWVAFASVPGIIFTK 180

QY 181 COKEDSVYVCGPYPRGWNFFHIMRNILGLVLPPLIMVICYSGILKTLRCRNEKKRHR 240

DB 181 COKEDSVYVCGPYPRGWNFFHIMRNILGLVLPPLIMVICYSGILKTLRCRNEKKRHR 240

QY 241 AVRVIPTIMIVFLFPTPNVIVLLNTFQEFFGLSNCESTSQLDQAVQVTTETLGMTHCCI 300

DB 241 AVRVIPTIMIVFLFPTPNVIVLLNTFQEFFGLSNCESTSQLDQAVQVTTETLGMTHCCI 300

QY 301 NPIIYAFVGEKFRRLSVFRRKHITRFRCKQCPVFYRETVDGVTSTNTPSTGEOEVSAGL 360

DB 301 NPIIYAFVGEKFRRLSVFRRKHITRFRCKQCPVFYRETVDGVTSTNTPSTGEOEVSAGL 360

## RESULT 6

ABB56340

ID ABB56340 standard; Protein; 360 AA.

XX AC ABB56340;

DT 18-FEB-2002 (first entry)

DE Non-endogenous human GPCR protein, SEQ ID NO: 473.

XX Human; G protein-coupled receptor; GPCR; non-endogenous; mutant;  
 KW constitutively activated GPCR; agonist; disease.

XX Homo sapiens.

OS Synthetic.

XX PN WO200177172-A2.

XX PD 18-OCT-2001.

XX PF 05-APR-2001; 2001WO-US11098.

XX PR 07-APR-2000; 2000US-195747P.

XX PA (AREN-) ARENA PHARM INC.

XX PI Lehmann-Bruinsma K, Liaw CW, Lin I;

XX DR WPI; 2001-648759/74.

DR N-PSDB; ABI97976.

PT Identifying agonists of G protein-coupled receptors (GPCRs) for use in  
 PT disease treatment, comprises contacting candidate compounds with  
 PT versions of GPCRs.

XX Claim 1; Page 274-275; 394pp; English.

XX The invention relates to G protein-coupled receptors (GPCRs) for which

CC the endogenous ligand has been identified. Non-endogenous  
 CC constitutively activated versions of known GPCRs are used in the  
 CC invention for the direct identification of candidate compounds as  
 CC receptor agonists, inverse agonists or partial agonists. Such  
 CC agonists are useful as therapeutic agents for diseases or disorders  
 CC associated with GPCRs. The present sequence is a non-endogenous  
 CC version of a known human GPCR.

XX Sequence 360 AA;

Query Match 99.7%; Score 1894; DB 22; Length 360;  
 Best Local Similarity 99.7%; Pred. No. 1.8e-210;  
 Matches 359; Conservative 0; Mismatches 1; Indels 0; Gaps 0;

Qy 1 MLSTSRFRIRNTNESGEEVTFDDYDYGAPCHKFDVKQIGQALLPPLSLVFIQFVGN 60  
 Db 1 MLSTSRFRIRNTNESGEEVTFDDYDYGAPCHKFDVKQIGQALLPPLSLVFIQFVGN 60

Qy 61 MLVVLILINCKKLCITDIYLLNLAISDLFLITPLWAHSAANEVFGNAMCKLFTGLY 120  
 Db 61 MLVVLILINCKKLCITDIYLLNLAISDLFLITPLWAHSAANEVFGNAMCKLFTGLY 120

Qy 121 HIGVGGIFFIILLIDRYLAIVHAFKARTVFGVTSVITWLVAVFASVPGIIFTK 180  
 Db 121 HIGVGGIFFIILLIDRYLAIVHAFKARTVFGVTSVITWLVAVFASVPGIIFTK 180

Qy 181 CQKEDSVVCGPYFPRGWNNEFTIMRNILGLVPLIMVICYSGILKTLRCRNEKKRHR 240  
 Db 181 CQKEDSVVCGPYFPRGWNNEFTIMRNILGLVPLIMVICYSGILKTLRCRNEKKRHR 240

Qy 241 AVRVIITMIVYFLWTPYNIIVLLNTFOEFGSLNCESTSLDQATQVETLGMTHCCI 300  
 Db 241 AVRVIITMIVYFLWTPYNIIVLLNTFOEFGSLNCESTSLDQATQVETLGMTHCCI 300

Qy 301 NPIIYAFVGEKFRRLYSVFERKHITKRCQCPVFVRETVDGVTSTNTSTGQEVSA 360  
 Db 301 NPIIYAFVGEKFRRLYSVFERKHITKRCQCPVFVRETVDGVTSTNTSTGQEVSA 360

RESULT 7

AAR79165  
 ID AAR79165 standard; Protein; 374 AA.

AC AAR79165;

XX 29-DEC-1995 (first entry)

XX Human monocyte chemoattractant protein-1 receptor MCP-1RA.

XX Monocyte chemoattractant protein-1 receptor; MCP-1R; chemokine.

XX Homo sapiens.

Key	Location/Qualifiers
FT Domain	49...70
FT	/label= transmembrane
FT Domain	80...700
FT	/label= transmembrane
FT Domain	115...136
FT	/label= transmembrane
FT Domain	154...178
FT	/label= transmembrane
FT Domain	204...231
FT	/label= transmembrane
FT Domain	244...268
FT	/label= transmembrane
FT Domain	295...313
FT	/label= transmembrane
FT Region	314...375
FT	/label= carboxyl tail
FT Domain	1..48
FT	/label= extracellular

PN WO9519436-A.  
 XX 20-JUL-1995.  
 XX 11-JAN-1995; 95WO-US00476.  
 XX 13-JAN-1994; 94US-0182962.  
 XX (REGC ) UNIV CALIFORNIA.  
 XX Charo I, Coughlin S;  
 PI WPI; 1995-263866/34.  
 XX N-PSDB; AAQ96297.

PT DNA encoding monocyte chemo-attractant protein-1 receptor - used partic.  
 PT for identifying antagonists and for treating diseases characterised by  
 PT monocytic infiltrates

XX Claim 2; Fig 1; 84pp; English.

CC To identify and clone new members of the chemokine receptor gene  
 CC family, degenerate oligo primers were designed corresp. to the  
 CC conserved sequences R79167 in the second and R79168 in the third  
 CC transmembrane domains of the MIP-1alpha/RANTES receptor, the IL-8  
 CC receptors and the HUMSTRS orphan receptor (GenBank Accession #M99293.  
 CC The degenerate oligo incorporating EcoRI and XhoI sites at their 5'  
 CC ends are Q96299 and Q96300. Amplification of cDNA derived from MM6  
 CC cells with the primers yielded a number of PCR products. One cDNA  
 CC appeared to encode a novel protein. To obtain a full-length version  
 CC of this clone, a MM6 cDNA library was constructed in pPROG and probed  
 CC with the PCR product. A 2.1 kb cDNA clone was obtd. Analysis of  
 CC additional clones in the MM6 cDNA library revealed a second  
 CC sequence that was identical to the 2.1 kb cDNA sequence first obtd.  
 CC from the 5' UTR through the putative seventh transmembrane domain  
 CC but contained a different cytoplasmic tail. The second sequence  
 CC appears to represent alternative splicing of the carboxyl-terminal  
 CC tail of the MCP-1R protein. The two sequences are denoted MCP-1RA  
 CC and MCP-1RB (see Q96297/R79165 & Q96298/R79166). Active mature  
 CC MCP-1RA has a predicted mol. wt. of about 42,000 daltons. MCP-1RB  
 CC has a mol. wt. of about 41,000 daltons.

XX Sequence 374 AA;

Query Match 86.9%; Score 1651.5; DB 16; Length 374;  
 Best Local Similarity 95.5%; Pred. No. 2.3e-182;  
 Matches 319; Conservative 3; Mismatches 5; Indels 7; Gaps 3;

Qy 1 MLSTSRFRIRNTNESGEEVTFDDYDYGAPCHKFDVKQIGQALLPPLSLVFIQFVGN 60  
 Db 1 MLSTSRFRIRNTNESGEEVTFDDYDYGAPCHKFDVKQIGQALLPPLSLVFIQFVGN 60

Qy 61 MLVVLILINCKKLCITDIYLLNLAISDLFLITPLWAHSAANEVFGNAMCKLFTGLY 120  
 Db 61 MLVVLILINCKKLCITDIYLLNLAISDLFLITPLWAHSAANEVFGNAMCKLFTGLY 120

Qy 121 HIGVGGIFFIILLIDRYLAIVHAFKARTVFGVTSVITWLVAVFASVPGIIFTK 180  
 Db 121 HIGVGGIFFIILLIDRYLAIVHAFKARTVFGVTSVITWLVAVFASVPGIIFTK 180

Qy 181 CQKEDSVVCGPYFPRGWNNEFTIMRNILGLVPLIMVICYSGILKTLRCRNEKKRHR 240  
 Db 181 CQKEDSVVCGPYFPRGWNNEFTIMRNILGLVPLIMVICYSGILKTLRCRNEKKRHR 240

Qy 241 AVRVIITMIVYFLWTPYNIIVLLNTFOEFGSLNCESTSLDQATQVETLGMTHCCI 300  
 Db 241 AVRVIITMIVYFLWTPYNIIVLLNTFOEFGSLNCESTSLDQATQVETLGMTHCCI 300

Qy 301 NPIIYAFVGEKFRRLYSVFERKHITKRCQCPV 334  
 Db 301 NPIIYAFVGEKFRRLYSVFERKHITKRCQCPV 334

## RESULT 8

AAG80107 standard; Protein; 374 AA.

XX  
AC AAG80107;  
XX  
DT 17-JAN-2002 (first entry)  
XX  
DE Human CCR2a protein.  
XX  
KW Chemokine; tumour diagnosis; colorectal; prostatic; organ rejection;  
KW inflammation; autoimmune disease; metastasis; bronchial asthma; lupus;  
KW chronic bowel inflammation; rheumatoid arthritis; cytostatic;  
KW antiinflammatory; antiasthmatic; immunosuppressive; dermatological;  
KW antirheumatic; antiarthritic.

XX Homo sapiens.

XX WO200172830-A2.

XX 04-OCT-2001.

XX 02-APR-2001; 2001WO-EP03708.

XX 31-MAR-2000; 2000DE-1016013.

XX (IPFP-) IPF PHARM GMBH.

XX (FORS/) FORSSMANN U.

XX Forssmann W, Adermann K, Heitland A, Spodsborg N;

XX WPI; 2001-626256/72.

XX Diagnostic agent containing two or more receptor-specific ligands,  
XX useful for detecting tumors, inflammation etc., also therapeutic use of  
XX ligand inhibitors

XX Disclosure; Page 9; 26pp; German.

XX This invention describes a novel diagnostic agent (A) comprising at least  
XX two different ligands (I) for receptors (II) that are implicated in  
XX disease. (A) are used for the diagnosis of tumors (especially colorectal  
XX or prostatic), organ rejection, inflammation and autoimmune diseases.  
XX Also inhibitors of (I) are used therapeutically against tumors (and their  
XX metastases), inflammation (particularly bronchial asthma or chronic bowel  
XX inflammation), or autoimmune diseases (rheumatoid arthritis or lupus),  
XX where the (cardio)vascular, lymphatic, respiratory, nervous, digestive,  
XX endocrine, motor or urogenital systems or skin are affected, and bone  
XX marrow diseases. The products of the invention are chemokine derivatives  
XX which have cytostatic, antiinflammatory, antiasthmatic,  
XX immunosuppressive, dermatological, antirheumatic, antiarthritic.  
XX Chemokines act on specific tumor and inflammatory cells through a  
XX constellation of chemokine receptors (CR), which control migration and  
XX proliferation of these cells. AAG80045-AAG80128 represent human chemokine  
XX fragments used to illustrate the method of the invention.

XX Sequence 374 AA;

Query Match 86.9%; Score 1651.5; DB 22; Length 374;  
Best Local Similarity 95.5%; Pred. No. 2.3e-182;  
Matches 319; Conservative 3; Mismatches 5; Indels 7; Gaps 3;

QY 1 MLSTSRSRIRNTNSESGETVTFDDYDYGAPCHKFDVKQIGAQLLPPLYSLVIFGFGVN 60  
DB 1 MLSTSRSRIRNTNSESGETVTFDDYDYGAPCHKFDVKQIGAQLLPPLYSLVIFGFGVN 60  
QY 61 MLVLLILNCKLKCLTDIYLLNLATSLLFLITLPLWAHSAANWVFGNACKLFTGLY 120  
DB 61 MLVLLILNCKLKCLTDIYLLNLATSLLFLITLPLWAHSAANWVFGNACKLFTGLY 120  
QY 121 HIGYFGGIFFIILLTDRLYLAIHVAFLKARTVTGVTSTWLVAVFASVPGIIFTK 180  
DB 121 HIGYFGGIFFIILLTDRLYLAIHVAFLKARTVTGVTSTWLVAVFASVPGIIFTK 180

QY 181 CQKEDSVYVCGPYFPRGWNNFHTIMRNILGLVLPPLIMVICYSGILKTLRCRNEKKRHR 240  
DB 181 CQKEDSVYVCGPYFPRGWNNFHTIMRNILGLVLPPLIMVICYSGILKTLRCRNEKKRHR 240  
QY 241 AVRVIETIMIVYFLEWTPYINIVILLNTFOEFGLSNCESTSOLDQATQVETILGMTHCCI 300  
DB 241 AVRVIETIMIVYFLEWTPYINIVILLNTFOEFGLSNCESTSOLDQATQVETILGMTHCCI 300  
QY 301 NPIIYAFVGEKFRYLSVFFERKHITKRCCKQCPV 334  
DB 301 NPIIYAFVGEKFRYLSVFFERKHITKRCCKQCPV 334

## RESULT 9

AAB46859

XX AAB46859 standard; Protein; 329 AA.

XX AAB46859;

XX 16-AUG-2001 (updated)

XX 02-AUG-2001 (updated)

XX 04-MAY-2001 (first entry)

XX Human MCP-1 receptor protein fragment.

XX HDGNR10; human; G-protein chemokine receptor; antiinflammatory;  
XX immunomodulatory; anticoagulant; antiallergic; immunosuppressive;  
XX cytostatic; antiparasitic; antipsoriatic; antirheumatic; antiarthritic;  
XX vasotropic; gene therapy; haematopoiesis; wound healing; coagulation;  
XX angiogenesis; solid tumour; infection; leukemia; growth factor activity;  
XX T-cell mediated autoimmune disease; psoriasis; allergy; atherogenesis;  
XX anaphylaxis; malignancy; inflammation; histamine; IgE; sfilicosis; shock;  
XX immunoglobulin E-mediated allergic reaction; rheumatoid arthritis;  
XX prostaglandin-independent fever; bone marrow failure; sarcoidosis;  
XX hyper-eosinophilic syndrome; vulnery.

XX Homo sapiens.

XX US2001000241-A1.

XX 12-APR-2001.

XX 29-NOV-2000; 2000US-0725285.

XX 06-JUN-1995; 95US-0466343.

XX 18-NOV-1998; 98US-0195662.

XX 25-JUN-1999; 99US-0339912.

XX (LIYY/) LI Y.

XX (RUBE/) RUBEN S M.

XX Li Y, Ruben SM;

XX WPI; 2001-226317/23.

XX New human G-protein chemokine receptor polypeptides and  
XX polynucleotides, useful for identifying (ant)agonists to the G-protein  
XX chemokine receptor

XX Disclosure; Page 16-17; 22pp; English.

XX This invention describes a novel receptor polypeptide (I) selected from  
XX (i) a fully defined 329 amino acid sequence (II) fully disclosed in the  
XX specification; and (ii) a polypeptide encoded by the cDNA contained in a  
XX plasmid, and fragments, analogs and derivatives of the polypeptide. The  
XX products of the invention have antiinflammatory, immunomodulatory,  
XX anticoagulant, antiallergic, immunosuppressive, vulnery, cytostatic,  
XX antiparasitic, antipsoriatic, antirheumatic, antiarthritic and vasotropic  
XX activity and can be used for gene therapy. The G-protein chemokine  
XX receptors, HDGNR10, (I) are useful for screening for compounds which  
XX activate or inhibit activation of (I). The products of the invention can  
XX also be used for stimulating haematopoiesis, wound healing, coagulation,

CC angiogenesis, treating solid tumours, chronic infections, leukemia, and  
CC T-cell mediated autoimmune diseases, parasitic infections, psoriasis, and  
CC stimulating growth factor activity. HDGN10 is useful for treating  
CC allergy, atherogenesis, anaphylaxis, malignancy, chronic and acute  
CC inflammation, histamine and immunoglobulin E (IgE)-mediated allergic  
CC reactions, prostaglandin-independent fever, bone marrow failure,  
CC silicosis, sarcoidosis, rheumatoid arthritis, shock and  
CC hyper-eosinophilic syndrome.  
CC (N.B. This record was resubmitted to correct errors in the keyword  
CC formatting).  
XX  
SQ

Sequence 329 AA;  
Query Match 77.5%; Score 1473; DB 22; Length 329;  
Best Local Similarity 90.5%; Pred. No. 9.3e-162;  
Matches 287; Conservative 3; Mismatches 5; Indels 22; Gaps 4;  
QY 18 EEVTFPDYDYGAPCHKFDVKQIGAQLLPPLYSLVFIFGVGNMLVLLINCKKLCIT 77  
DB 1 EEVTFPDYDYGAPCHKFDVKQIGAQLLPPLYSLVFIFGVGNMLVLLINCKKLCIT 60  
QY 78 DIYLLNLAIISDLLFLITPLMAHSAANEWFVGNAMCKLFTGLYHIGYFGGIFITLITID 137  
DB 61 DIYLLNLAIISDLLFLITPLMAHSAANEWFVGNAMCKLFTGLYHI----- 105  
QY 138 RYLAIVHAVFALKARTVTFGVVSVITWLVAVFASVPGIIFTKCKEDSVVCGPYPRG 197  
DB 106 RYLAIVHAVFALKARTVTFGVVSVITWLVAVFASVPGIIFTKCKEDSVVCGPYPRG 165  
QY 198 WNNFHTIMRNILGLVPLLMVICYSGILKTLRCRNEKKRHRVAVRVIITMIVVFLFWT 257  
DB 166 WNNFHTIMRNILGLVPLLMVICYSGILKTLRCRNEKKRHRVAVRVIITMIVVFLFWT 225  
QY 258 PYNIVILLNTFOEFFGLSNCESTSQLDQATQVTTGLMTHCCINPIIYAFVGEKFRYLS 317  
DB 226 PYNIVILLNTFOEFFGLSNCESTSQLDQATQVTTGLMTHCCINPIIYAFVGEKFRYLS 282  
QY 318 VFFRKHIHTRKCKQCPV 334  
DB 283 LF---HIALG-CRIAPL 295

RESULT 10  
AAW54037  
ID AAW54037 standard; Protein; 354 AA.  
XX  
AC AAW54037;  
XX  
PT 06-AUG-1998 (first entry)  
XX  
DE Mouse CC-CR5 protein.  
XX  
XX CC-CR5; chemokine receptor; mouse; human; transgenic mouse;  
KW HIV infection; T-cell mediated inflammation.  
XX  
OS Mus sp.  
XX  
PN EP834564-A2.  
XX  
PD 08-APR-1998.  
XX  
PF 03-OCT-1997; 97EP-0307823.  
XX  
XX 03-OCT-1996; 96US-0724984.  
PR (SMIK ) SMITHKLINE BEECHAM CORP.  
XX  
PA Bergsma DJ, Brawner ME, Shabon U;  
XX  
PI WPI; 1998-195463/18.  
XX  
DR N-PSDB; AAV23989.  
XX  
PT New isolated mouse chemokine receptor, CC-CR5 - used to develop

PT products for the study, diagnosis and treatment of HIV infection or  
PT T-cell mediated inflammation  
XX  
PS Claim 11; Fig 1; 27pp; English.  
XX  
CC This sequence is the mouse CC-CR5 protein, is encoded by the DNA of the  
CC invention. CC-CR5 is a chemokine receptor. Cells transfected with the  
CC DNA can be cultivated and the expression product harvested. The DNA can  
CC be knocked out and replaced with the human CC-CR5 gene, creating  
CC transgenic mice which can be used in the study of HIV infection or T-cell  
CC mediated inflammation. Transgenic mice could also be used to screen for  
CC human CC-CR5 agonists or antagonists.  
XX  
SQ

Sequence 354 AA;  
Query Match 72.9%; Score 1386; DB 19; Length 354;  
Best Local Similarity 74.0%; Pred. No. 1.2e-151;  
Matches 259; Conservative 33; Mismatches 52; Indels 6; Gaps 2;  
QY 17 GEEVTFPDYDYG--APCHKFDVKQIGAQLLPPLYSLVFIFGVGNMLVLLINCKKLCIT 74  
DB 5 GSVPTIYDIDYGSAPCQKINVKQIAAQLLPPLYSLVFIFGVGNMLVLLINCKKLCIT 64  
QY 75 CLTDIYLLNLAIISDLLFLITPLMAHSAANEWFVGNAMCKLFTGLYHIGYFGGIFITLIT 134  
DB 65 SVTDIYLLNLAIISDLLFLITPLFWAHYAANEWFVGNAMCKLFTGLYHIGYFGGIFITLIT 124  
QY 135 TIDRYLAIVHAVFALKARTVTFGVVSVITWLVAVFASVPGIIFTKCKEDSVVCGPYPRG 194  
DB 125 TIDRYLAIVHAVFALKARTVTFGVVSVITWLVAVFASVPGIIFTKCKEDSVVCGPYPRG 184  
QY 195 PRG----WNNFHTIMRNILGLVPLLMVICYSGILKTLRCRNEKKRHRVAVRVIITM 250  
DB 185 PHTQYHFNKSFQTLKMWILSLILPLLMVICYSGILKTLRCRNEKKRHRVAVRVIITM 244  
QY 251 VYFLEWTPYNIIVILLNTFOEFFGLSNCESTSQLDQATQVTTGLMTHCCINPIIYAFVGE 310  
DB 245 VYFLEWTPYNIIVILLNTFOEFFGLSNCESTSQLDQATQVTTGLMTHCCINPIIYAFVGE 304  
QY 311 KFRYSLSVFFRKHIHTRKCKQCPVRETVDGVTSTNTPTSGEQEVSAGL 360  
DB 305 KFRYSLSVFFRKHIHTRKCKQCPVRETVDGVTSTNTPTSGEQEVSAGL 354

RESULT 11  
AAG79089  
ID AAG79089 standard; Protein; 352 AA.  
XX  
AC AAG79089;  
XX  
DT 10-DEC-2001 (first entry)  
XX  
DE Amino acid sequence of human CCR5 protein.  
XX  
XX Human; receptor; DC-SIGN; dendritic cell; T lymphocyte; HIV;  
KW gp120; C-type lectin; ICAM3; HIV entry; T cell; macrophage;  
KW HIV infection; CCR5.  
XX  
OS Homo sapiens.  
XX  
PN WO200164752-A2.  
XX  
PD 07-SEP-2001.  
XX  
PF 28-FEB-2001; 2001WO-US06322.  
XX  
XX 02-MAR-2000; 2000US-0517605.  
PR (UYNY ) UNIV NEW YORK STATE.  
XX  
PA (UYNI-) UNIV NIJMEGEN.  
XX  
PI Littman DR, Kwon D, Van Kooyk Y, Geijtenbeek T;  
XX

DR WPI; 2001-602565/68.  
 XX  
 CC An antibody for the treatment or prevention of HIV-infection comprises  
 PT a gp120 portion which binds to DC-SIGN or is exposed upon gp120 binding  
 PT of DC-SIGN due to concomitant conformational change -  
 XX  
 PS Disclosure; Page 118-119; 131pp; English.  
 XX  
 CC The specification describes an antibody which is specific for an  
 CC antigenic fragment of gp120. This antigenic fragment binds to DC-SIGN  
 CC or is exposed upon gp120 binding of DC-SIGN due to concomitant  
 CC conformational change. DC-SIGN is a receptor that is specifically  
 CC expressed on dendritic cells and facilitates infection of T lymphocytes  
 CC with HIV. DC-SIGN is identical to a HIV-1 gp120-binding C-type lectin.  
 CC DC-SIGN binds ICAM-3 (which is expressed constitutively on T lymphocytes)  
 CC with high affinity. The antibody of the invention inhibits the trans  
 CC enhancement of HIV entry into a T cell or macrophage facilitated by  
 CC dendritic cells. The antibody is useful to treat or prevent HIV  
 CC infection. The present sequence represents a human CCR5 protein,  
 CC which is a translocation promoting agent that interacts with CD4.  
 CC This receptor functions in HIV-1 entry into cells.

SQ Sequence 352 AA;

Query Match 72.2%; Score 1371; DB 22; Length 352;

Best Local Similarity 76.2%; Pred. No. 6.6e-150;

Matches 259; Conservative 30; Mismatches 47; Indels 4; Gaps 1;

QY 25 DYDYGAPCHKFDVKQIGAOQLPPLYSILVIFGFGVGNMVLVILINCKKLCTDIYLLN 84  
 DB 13 DYTSEPCQKINVKQIAARLLPPLYSILVIFGFGVGNMVLVILINCKKLCTDIYLLN 72  
 QY 85 AYSDDLFLITLPWAHSAANWVFGNAMCKLFTGLYHIGYFGGIFILLTIDRYLAIVH 144  
 DB 73 AISDLFLITVPWAHAAQWDFGNTMCQLLTGLYFGGIFILLTIDRYLAIVH 132  
 QY 145 AVFALKARTVTEGVVTSVITLWVAVFASVPGIIFTCOKEDSVVCGPYFP---RGWNN 200  
 DB 133 AVFALKARTVTEGVVTSVITWVAVFASVPGIIFTRSQEGHLHYTCSSHPFYSOYQF 192  
 QY 201 FHTIMNIGLVLPLIMVICYSGILKTLRLCRNEKKRHRVAVIFTIMIVYFLFWT 260  
 DB 193 FQTLKIVILGLVPLVIMVICYSGILKTLRLCRNEKKRHRVAVIFTIMIVYFLFWAPN 252  
 QY 261 IVLLNTFOEFFGLSNCESTSQLDQATQVTELTGMTHCCINPIIYAFVGEKFRYLSVF 320  
 DB 253 IVLLNTFOEFFGLNCCSSNRLDQAMQVTELTGMTHCCINPIIYAFVGEKFRYLLVF 312  
 321 RKHITKRCKQCPVYRETVDGVTSTNTPTSGEQEVSAGL 360  
 313 QKHIAKHFCKCCSIFQOEAPERASSVYTRSTGQEISVGL 352

RESULT 12  
 AAW27407

ID AAW27407 standard; Protein; 352 AA.

XX AAW27407;

AC AAW27407;

DT 14-APR-1998 (first entry)

XX Human CCR5.

KW Human Cys-Cys chemokine receptor 5; CCR5;

KW human immunodeficiency virus; type 1; type 2; HIV-1; HIV-2;

KW diagnosis; treatment; prevention;

KW inflammatory disease; rheumatoid arthritis; glomerulonephritis;

KW asthma; idiopathic pulmonary fibrosis; psoriasis; viral infection;

KW cancer; atherosclerosis; autoimmune disorder.

OS Homo sapiens.

XX WO9732019-A2.

PN

XX 04-SEP-1997.  
 PD  
 XX 28-FEB-1997; 97WO-BE00023.  
 PF  
 XX 06-AUG-1996; 96EP-0870102.  
 PR  
 XX 01-MAR-1996; 96EP-0870021.  
 PR  
 XX (EURO-) EUROSREEN SA.  
 PA  
 XX Libert F, Parmentier M, Samson M, Vassart G;  
 PI  
 XX WPI; 1997-479829/44.  
 DR  
 XX N-PSDB; AAT90117.  
 DR  
 PT Active and inactive forms of human CC chemokine receptor CCR-5  
 PT useful to diagnose, prevent and/or treat inflammatory disorders,  
 PT autoimmune disease and viral infection  
 XX  
 PS Claim 4; Fig 1b-c; 94pp; English.

XX The present sequence is human CC (Cys-Cys) chemokine receptor  
 CC 5 (CCR5), which is stimulated by MIP-1 alpha, MIP-1 beta or RANTES  
 CC chemokines, but not by monocyte chemoattractant protein 1 (MCP-1),  
 CC MCP-2, MCP-3, interleukin-8 (IL-8) or growth related gene product  
 CC alpha (GRO alpha) chemokines. Active CCR-5 is also a receptor of  
 CC human immunodeficiency virus type 1 or type 2 (HIV-1 or HIV-2).  
 CC CCR5 or its cDNA can be used to diagnose, treat and/or prevent  
 CC inflammatory diseases, e.g. rheumatoid arthritis,  
 CC glomerulonephritis, asthma, idiopathic pulmonary fibrosis and  
 CC psoriasis, viral infections, especially HIV-1 or HIV-2 infection,  
 CC cancer, atherosclerosis and autoimmune disorders.

XX Sequence 352 AA;

Query Match 71.8%; Score 1364; DB 18; Length 352;

Best Local Similarity 75.5%; Pred. No. 4.3e-149;

Matches 259; Conservative 32; Mismatches 46; Indels 6; Gaps 2;

QY 24 FDYDY--GAPCHKFDVKQIGAOQLPPLYSILVIFGFGVGNMVLVILINCKKLCTDIYLL 81  
 DB 10 YDINYTSEPCQKINVKQIAARLLPPLYSILVIFGFGVGNMVLVILINCKRLKSM 69  
 QY 82 LNLAISDDLFLITLPWAHSAANWVFGNAMCKLFTGLYHIGYFGGIFILLTIDRYLA 141  
 DB 70 LNLAISDDLFLITVPWAHAAQWDFGNTMCQLLTGLYFGGIFILLTIDRYLA 129  
 QY 142 IVHAFVALKARTVTEGVVTSVITLWVAVFASVPGIIFTCOKEDSVVCGPYFP---RG 197  
 DB 130 VVHAFVALKARTVTEGVVTSVITWVAVFASVPGIIFTRSQEGHLHYTCSSHPFYSOYQ 189  
 QY 198 WNFHTIMRNILGLVPLIMVICYSGILKTLRLCRNEKKRHRVAVIFTIMIVYFLFWT 257  
 DB 190 WNFHTIMRNILGLVPLIMVICYSGILKTLRLCRNEKKRHRVAVIFTIMIVYFLFWA 249  
 QY 258 PYNIVILLNTFOEFFGLSNCESTSQLDQATQVTELTGMTHCCINPIIYAFVGEKFRYLS 317  
 DB 250 PYNIVILLNTFOEFFGLNCCSSNRLDQAMQVTELTGMTHCCINPIIYAFVGEKFRYLL 309  
 QY 318 VFERKHITKRCKQCPVYRETVDGVTSTNTPTSGEQEVSAGL 360  
 DB 310 VFERKHITKRCKQCPVYRETVDGVTSTNTPTSGEQEVSAGL 352

RESULT 13

AAW27123

ID AAW27123 standard; Protein; 352 AA.

XX AAW27123;

AC AAW27123;

DT 14-DEC-1997 (first entry)

XX Human chemokine receptor 88C.



RESULT 14

AA  
AC  
AAW27125:

14-DEC-1997 (first entry)

XX  
DE  
Macaque chemokine receptor

chemokine receptor 88C: atherosclerosis

KW tumour; asthma; viral infection; AIDS; inflammation;  
KW autoimmune disease; therapy; diagnosis; leukocyte trafficking;  
KW autoimmunity; ligand; modulator; antibody.

XX  
OS Macaca sp.

WO9722698-A2.

XX : 26-JUN-1997.  
PDXX  
PF  
20-DEC-1996: 96WO-US20759.XX  
PB  
07-JUN-1996. 96US-0661393.

PR 20-DEC-1993; 9305-03/3987;  
yy

PA (ICOS-) ICOS CORP.

PI Gray PW, Raport CJ, Schweickart VL,

DR WPI; 1997-341689/31.  
N-DCDR: 3AT95163

[illegible]

to modulate leukocyte trafficking, e.g. for treatment of inflammation, tumours, viral infections, autoimmune diseases, etc.

XX  
21-58 26. Page 57-58. 65nn. English.

XX This polypeptide sequence comprises macaque chemokine receptor 88C,  
CC a G protein coupled receptor that is involved in leukocyte  
CC trafficking. The amino sequence was deduced from a 88C DNA  
CC sequence.

CC (AAT85163) isolated by PCR amplification. It shows 97% identity to 98C receptors and their polypeptide fragments

CC can be produced in transformed host cells. The receptors, peptides

CC domains, and anti-receptor antibodies can be used to modulate

CC are potentially potentially useful in the treatment of

infection, AIDS, inflammatory conditions, pathological immune

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
84

2

CC that produces an antibody that specifically binds to macaque 88C is  
 CC claimed.

XX Sequence 352 AA;

Query Match 71.8%; Score 1364; DB 18; Length 352;  
 Best Local Similarity 75.5%; Pred. No. 4.3e-149;  
 Matches 260; Conservative 31; Mismatches 46; Indels 6; Gaps 2;

QY 24 FDYDY--GAPCHKFDVKQIGAOQLPPLYSLVFIFGVGNMVLVILINCKKLKCLTDIYL 81  
 Db 10 YDIDYTTSEPCQKINVKQIAARLLPPLYSLVFIFGVGNMVLVILINCKKLKCLTDIYL 69

QY 82 LNLAISSDLLFLITLPLWAHSAANEWVFGNACKLFTGLYHIGYFGGIFFTLLTIDRYLA 141  
 Db 70 LNLAISSDLLFLITLPLWFAHAAQWDFGNTMCQLLTGLYFIFGFGIFFTLLTIDRYLA 129

QY 142 IVHAVFALKARTVTGVTSTWLVAVFASVPGIIFTRKQKEDSVYVCGPYFP---RG 197  
 Db 130 IVHAVFALKARTVTGVTSTWLVAVFASVPGIIFTRKQKEDSVYVCGPYFP---RG 189

QY 198 WNNFHTIMRNILGLVPLIMVICYSGILKTLRCRNEKRRHRAVRVIFTIMIVYFLFWT 257  
 Db 190 WNFQTLKAVILGLVPLIMVICYSGILKTLRCRNEKRRHRAVRVIFTIMIVYFLFWT 249

QY 258 PYNIVILLNTFOEFGFLSNCESTSDQATQVTELTGMTHCCINPIIYAFVGEKFRYLS 317  
 Db 250 PYNIVILLNTFOEFGFLSNCESTSDQATQVTELTGMTHCCINPIIYAFVGEKFRYLS 309

QY 318 VFFRKHITKRCCKOCPVYRETVDGVTSTNTPSTGEQEVSAGL 360  
 Db 310 VFFOKHIAKRCCKCSIFQOEAPERASSVYTRSTGEQEISVGL 352

## RESULT 15

AAW23835

ID AAW23835 standard; Protein; 352 AA.

XX AAW23835;

XX 08-JUN-1998 (first entry)

XX Human CC chemokine receptor 5 (CCR5).

DE CC chemokine receptor 5; CCR5; G-protein coupled receptor;  
 KW human immunodeficiency virus; HIV; CD4; AIDS; therapy;  
 KW transgenic animal.

XX Homo sapiens.

Key Location/Qualifiers  
 Domain 29..55  
 /label= I  
 Region 109..120  
 /note= "transmembrane domain"  
 Domain 104..126  
 /label= III  
 /note= "extracellular loop-1 (Claim 19)"  
 Domain 143..171  
 /label= IV  
 /note= "transmembrane domain"  
 Region 187..210  
 /label= V  
 /note= "transmembrane domain"  
 Domain 194..219  
 /label= VI  
 /note= "transmembrane domain"  
 Region 238..258  
 /label= VII  
 /note= "transmembrane domain"  
 Region 261..276  
 /note= "transmembrane domain"  
 Domain 277..300  
 /note= "extracellular loop-3 (Claim 19)"

FT /label= VII  
 FT /note= "transmembrane domain"

XX WO9745543-A2.

XX 04-DEC-1997.

XX 28-MAY-1997; 97WO-US09586.

XX 28-MAY-1996; 96US-0018508.

XX (USSH ) US DEPT HEALTH & HUMAN SERVICES.

XX Alkhatib G, Berger EA, Broder CC, Combadiere C;  
 PI Feng Y, Kennedy PE, Murphy PM;

XX WPI; 1998-032650/03.

XX N-PSDB; AAT76920.

XX CC chemokine receptor 5 polypeptide - used to inhibit membrane  
 fusion between HIV and a target cell

XX Claim 68; Fig 1C; 70pp; English.

XX This protein sequence comprises of a novel human macrophage-selective  
 CC chemokine receptor that has been designated CCR5. The sequence  
 was deduced from an isolated cDNA clone (see AAT76920). An Alai27Leu  
 variant (see W38340 of CCR5 was also identified. The susceptibility  
 of human macrophages to HIV infection depends on cell surface  
 expression of CD4 and CCR5. CCR5 is a member of the 7-transmembrane  
 superfamily of G-protein coupled cell surface molecules. It plays  
 an essential role in the membrane fusion step of infection by some  
 CC HIV isolates. The establishment of stable, non-human cell lines  
 CC and transgenic mammals having cells that coexpress human CD4 and  
 CCR5 provides valuable tools for research of HIV infection.  
 CC Antibodies that bind to CCR5, CCR5 variants, and CCR5-binding  
 CC agents capable of blocking membrane fusion between HIV and target  
 CC cells represent potential anti-HIV therapeutics for macrophage  
 CC tropic strains of HIV.

XX Sequence 352 AA;

Query Match

Best Local Similarity 71.8%; Score 1364; DB 19; Length 352;  
 Matches 259; Conservative 32; Mismatches 46; Indels 6; Gaps 2;

QY 24 FDYDY--GAPCHKFDVKQIGAOQLPPLYSLVFIFGVGNMVLVILINCKKLKCLTDIYL 81  
 Db 10 YDIDYTTSEPCQKINVKQIAARLLPPLYSLVFIFGVGNMVLVILINCKKLKCLTDIYL 69

QY 82 LNLAISSDLLFLITLPLWAHSAANEWVFGNACKLFTGLYHIGYFGGIFFTLLTIDRYLA 141  
 Db 70 LNLAISSDLLFLITLPLWFAHAAQWDFGNTMCQLLTGLYFIFGFGIFFTLLTIDRYLA 129

QY 142 IVHAVFALKARTVTGVTSTWLVAVFASVPGIIFTRKQKEDSVYVCGPYFP---RG 197  
 Db 130 IVHAVFALKARTVTGVTSTWLVAVFASVPGIIFTRKQKEDSVYVCGPYFP---RG 189

QY 198 WNNFHTIMRNILGLVPLIMVICYSGILKTLRCRNEKRRHRAVRVIFTIMIVYFLFWT 257  
 Db 190 WNFQTLKAVILGLVPLIMVICYSGILKTLRCRNEKRRHRAVRVIFTIMIVYFLFWT 249

QY 258 PYNIVILLNTFOEFGFLSNCESTSDQATQVTELTGMTHCCINPIIYAFVGEKFRYLS 317  
 Db 250 PYNIVILLNTFOEFGFLSNCESTSDQATQVTELTGMTHCCINPIIYAFVGEKFRYLS 309

QY 318 VFFRKHITKRCCKOCPVYRETVDGVTSTNTPSTGEQEVSAGL 360  
 Db 310 VFFOKHIAKRCCKCSIFQOEAPERASSVYTRSTGEQEISVGL 352

Search completed: May 19, 2003, 16:45:37  
 Job time : 54.9605 secs

us-09-625-573-4.rag

Tue May 20 09:36:44 2003

